

Media Coverage of Roselle Muffin

Developed by
Amity Food and Agriculture Foundation
Amity University
Uttar Pradesh, Noida



A panel of 30 people tested a total of 17 recipe variants in the muffin study. The aim was to find the best version of Roselle muffins in relation to 3 important qualities: Taste, nutrition and shelf life. Illustration photo: Pexels

Muffins that could be good for your health

Imagine sinking your teeth into a tender muffin that tastes good and is chock full of vitamins, antioxidants and other natural ingredients that is also good for your health, too.

By [Selvi Normannsen](#) - Published 07.03.2023

Love muffins? We're talking about a tasty, fluffy muffin that has no artificial additives and that simultaneously contains lots of beneficial nutrients. As remarkable as it might sound, a recently published study led describes the development of just such a new muffin in the journal *Foods*.

Super muffins? Research adds health benefits with hibiscus extract

08 Mar 2023 -- Researchers developed a muffin recipe containing an optimal concentration of Roselle calyx extract from the husks of *Hibiscus sabdariffa*. Dubbed the Roselle muffin, a panel positively tested the product's taste and texture. The muffin contained high levels of antioxidants, anthocyanin, phenolic and ascorbic acid, which the researchers note are linked to various health benefits.

NutritionInsight sits down with the study's authors, Eva Falch, associate professor in food science at the Norwegian University of Science and Technology, and Nutan Kaushik, professor at the Amity University Uttar Pradesh in India.

"We hope that this study will widen the use of natural ingredients in food for health, taste, and preservation purposes," notes Falch.

"At the same time, this study increases the knowledge about underutilized food resources that can be upgraded as future food ingredients, thereby contributing to food and nutritional security. Many of our global future food sources are scarce and as researchers, I think we need to engage also in the formulation of new healthy products."

The authors note that the study's outcomes can be utilized to develop foods containing Roselle calyx extract.

The study is part of Re-FOOD, a Norwegian-Indian collaborative project that focuses on using and enhancing the value of the left-over raw materials from food processing. The results have been published in *Foods*.



The researchers created a muffin containing Roselle calyx extract that had nutritional benefits.

Commercial applications

Falch explains that business development is needed to upgrade Roselle extracts into food ingredients, as these are not commercially available.

"One way of marketing the product's health benefits is using health and nutrition claims on the front-of-pack label. Different authorities regulate these depending on the targeted market and what scientific substantiation is needed."

Kaushik suggests that food manufacturers target health-conscious consumers with the product. "The plus point is that the product contains no artificial ingredients."

The Roselle muffin has a shelf life of six days without preservatives.

Health benefits

Roselle calyx extract contains [bioactive compounds](#), such as polyphenols, flavonoids, betaine and hibiscus acid.

However, the researchers note that antioxidant properties decreased after baking. The muffin batter had an antioxidant activity of 27.3%, while that of the baked Roselle muffin was 12.53%.

"Roselle extracts also contain compounds that are important in a healthy diet, such as the cocktail of antioxidants that need to be further explored in some markets to use health claims," adds Falch.

"As an ingredient, the Roselle calyx extract can provide different positive properties to a product such as bioactive, natural preservation and natural color."

Optimizing muffin mix

The product was developed using different formulations in varying proportions resulting from Response Surface Methodology that predicted formulations with the most favorable outcomes. The researchers replaced milk in vanilla muffins with varying concentrations of roselle extract and added different citric acid and sodium bicarbonate levels.

Thirty people tested the resulting 17 recipes. The optimal mix contained 45.37 ml of the extract volume, 1.11 grams of citric acid and 1.16 grams of sodium bicarbonate, which scored 8.31 for texture and 8.30 for overall acceptability.

The researchers measured the participants' responses to the muffins' color, appearance, aroma, texture, fluffiness, elasticity, freshness and chewing resistance.

Compared to a control muffin, the optimized Roselle muffin had higher acceptability scores for body, texture, taste and flavor. However, some taste panelists noted the muffin had a slightly sour taste.

Follow-up research

Falch notes that the study has demonstrated a method to scientifically optimize the functional and sensory properties of a new ingredient in a product, with muffins as a model product. "The nutritional composition of the muffin recipe can be improved by reducing the sugar and fat content."

Could a muffin be the next functional food?

[re-FOOD](#)

Friday, 10 March, 2023

[Share](#) [Tweet](#) [Share](#) [Email](#)



Imagine sinking your teeth into a tender muffin that tastes good and is chock full of vitamins, antioxidants and other natural ingredients that are good for your health, too. Researchers are testing possible varieties.

This super muffin has been named Roselle, because it contains calyx extract from the tropical plant *Hibiscus sabdariffa*, often referred to by the same name.

Calyx extract from the hibiscus flowers contains many bioactive compounds, such as polyphenols, flavonoids, betaine and hibiscus acid, all of which can be used to develop functional products that may offer positive health effects. The study also shows that Roselle muffins maintain freshness for up to six days at room temperature.

The study is part of [Re-FOOD](#), a Norwegian-Indian collaborative project that focuses on using and enhancing the value of raw materials of food processing.

Professor Nutan Kaushik at [Amity University](#) in the state of Uttar Pradesh, India, points out the health benefits of antioxidants, especially when it comes to neutralising free radicals, which can be an important contributing factor to serious illnesses.

<https://www.foodprocessing.com.au/content/ingredients/news/could-a-muffin-be-the-next-functional-food-1101340722>



(Credit: Openverse)

FOOD NEWS, HEALTH & MEDICAL NEWS

Adding this natural ingredient creates a healthier and longer-lasting muffin

Super muffins? Research adds health benefits with hibiscus extract

08 Mar 2023 --- Researchers developed a muffin recipe containing an optimal concentration of Roselle calyx extract from the husks of *Hibiscus sabdariffa*. Dubbed the Roselle muffin, a panel positively tested the product's taste and texture. The muffin contained high levels of antioxidants, anthocyanin, phenolic and ascorbic acid, which the researchers note are linked to various health benefits.

NutritionInsight sits down with the study's authors, Eva Falch, associate professor in food science at the Norwegian University of Science and Technology, and Nutan Kaushik, professor at the Amity University Uttar Pradesh in India.

"We hope that this study will widen the use of natural ingredients in food for health, taste, and preservation purposes," notes Falch.

"At the same time, this study increases the knowledge about underutilized food resources that can be upgraded as future food ingredients, thereby contributing to food and nutritional security. Many of our global future food sources are scarce and as researchers, I think we need to engage also in the formulation of new healthy products."

The authors note that the study's outcomes can be utilized to develop foods containing Roselle calyx extract.

The study is part of Re-FOOD, a Norwegian-Indian collaborative project that focuses on using and enhancing the value of the left-over raw materials from food processing. The results have been published in [Foods](#).



The researchers created a muffin containing Roselle calyx extract that had nutritional benefits.

Commercial applications

Falch explains that business development is needed to upgrade Roselle extracts into food ingredients, as these are not commercially available.

"One way of marketing the product's health benefits is using health and nutrition claims on the front-of-pack label. Different authorities regulate these depending on the targeted market and what scientific substantiation is needed."

Kaushik suggests that food manufacturers target health-conscious consumers with the product. "The plus point is that the product contains no artificial ingredients."

The Roselle muffin has a shelf life of six days without preservatives.

Health benefits

Roselle calyx extract contains [bioactive compounds](#), such as polyphenols, flavonoids, betaine and hibiscus acid.

However, the researchers note that antioxidant properties decreased after baking. The muffin batter had an antioxidant activity of 27.3%, while that of the baked Roselle muffin was 12.53%.

"Roselle extracts also contain compounds that are important in a healthy diet, such as the cocktail of antioxidants that need to be further explored in some markets to use health claims," adds Falch.

"As an ingredient, the Roselle calyx extract can provide different positive properties to a product such as bioactive, natural preservation and natural color."

Optimizing muffin mix

The product was developed using different formulations in varying proportions resulting from Response Surface Methodology that predicted formulations with the most favorable outcomes. The researchers replaced milk in vanilla muffins with varying concentrations of roselle extract and added different citric acid and sodium bicarbonate levels.

Thirty people tested the resulting 17 recipes. The optimal mix contained 45.37 ml of the extract volume, 1.11 grams of citric acid and 1.16 grams of sodium bicarbonate, which scored 8.31 for texture and 8.30 for overall acceptability.

The researchers measured the participants' responses to the muffins' color, appearance, aroma, texture, fluffiness, elasticity, freshness and chewing resistance.

Compared to a control muffin, the optimized Roselle muffin had higher acceptability scores for body, texture, taste and flavor. However, some taste panelists noted the muffin had a slightly sour taste.

Follow-up research

Falch notes that the study has demonstrated a method to scientifically optimize the functional and sensory properties of a new ingredient in a product, with muffins as a model product. "The nutritional composition of the muffin recipe can be improved by reducing the sugar and fat content."

Media Coverage of Roselle Muffin

The authors conclude that the ingredient should be tested in products with a better nutritional profile, such as whole grain products.

"A follow-up in this study will be to test the extract in different food systems and nutritionally optimized foods such as a Roselle muffin with minimal sugar, salt and saturated fat and maybe also more fibers," adds Falch. "It can then become a natural daily food choice replacing supplements."

"Another follow-up will be to document further the health benefits of the Roselle calyx extract and the effect in the food product used."

By Jolanda van Hal

To contact our editorial team please email us at editorial@cnsmedia.com



Roselle calyx extract contains bioactive compounds, such as antioxidants, flavonoids, betaine and hibiscus acid.



A panel of 30 people tested a total of 17 recipe variants in the muffin study. The aim was to find the best version of Roselle muffins in relation to 3 important qualities: Taste, nutrition and shelf life. Illustration photo: Pexels

Muffins that could be good for your health

Imagine sinking your teeth into a tender muffin that tastes good and is chock full of vitamins, antioxidants and other natural ingredients that is also good for your health, too.


By [Sølvi Normannsen](#) - Published 07.03.2023

Love muffins? We're talking about a tasty, fluffy muffin that has no artificial additives and that simultaneously contains lots of beneficial nutrients. As remarkable as it might sound, a recently published study led describes the development of just such a new muffin in the journal *Foods*.

From muffins to functional food

The new super muffin has been named Roselle, because it contains calyx extract from the tropical plant *Hibiscus sabdariffa*, which is often referred to by the same name.

Figure 1. Images of Roselle muffin obtained with different formulations combinations.



Here's a look at the different formulations that researchers tested. Photo: Screenshot from Food article

Hibiscus flowers have a beautiful red colour, and the calyx extract contains many valuable bioactive compounds, such as polyphenols, flavonoids, betaine and hibiscus acid. All of these components can be used to develop new products and functional food that may offer positive health effects, possibly also reducing the risk of chronic diseases.

The study also shows that Roselle muffins maintain their freshness for up to six days at room temperature, but with so many beneficial qualities, chances are they will be long gone by then.

Health benefits

The study is part of Re-FOOD (see factbox), a Norwegian-Indian collaborative project focusing on using and enhancing the value of rest raw materials of food processing.

RE-Food (2017-2020)

- The muffin-study is part of the Re-FOOD project, and INTPART-project funded by the Research Council of Norway.
- INTPART is an abbreviation for International Partnerships for Excellent Education, Research and Innovation
- The project builds collaboration between Norwegian and Indian research, education and innovation.
- The aim is to achieve the highest possible utilization of resources in the food value chain.
- RE-FOOD also focuses on the importance of natural plant extracts for colour, durability and health effects in the food of the future.

Professor Nutan Kaushik at Amity University in the state of Uttar Pradesh, India, points out the health benefits of antioxidants, especially when it comes to neutralizing free radicals, which can be an important contributing factor to serious illnesses.

Prefer natural food colouring

Roselle is also rich in anthocyanins. This is a dye that can be dissolved in water and yields a red, pink, purple, blue, blue violet or violet colour. The colour of food has a strong influence on how appetizing we think it is.

Artificial dyes are often clearer and more intense than natural substances. But the demand for natural dyes is increasing because of the concern that artificial dyes are harmful to our health.

Tasty, nutritional and long shelf-life

In the muffin study, 30 people tested a total of 17 recipe variants.

The aim was to find the best version of Roselle muffins relative to three important qualities:

- The muffin should have a taste and texture that people like
- The muffin should have valuable nutritional properties
- The muffin should have the best possible shelf life without preservatives

Aim — the best possible response

The researchers measured and surveyed the test subjects' response to the pastry's colour, appearance, aroma, texture, fluffiness, elasticity, freshness and chewing resistance. The researchers used the Response Surface Methodology, to model and analyse how the test subjects' reactions were affected by various factors. The aim of the method is to achieve the most positive response to what is tested.

Natural food colouring — good for our cells

In terms of nutrition, the results suggest that the hibiscus extract provides muffins with high levels of antioxidants like phenolics.

These are substances believed to be able to prevent processes in the body that can damage cell membranes or the genetic material in our cells. There are several thousand different phenols, most of which are found in fruit, juice, wine, cocoa, tea, coffee, vegetables, flax seed, whole grains and legumes.



Hibiscus sabdariffa is frequently used in teas and juices. Photo: Colourbox.

The muffin study

- Title: "Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology".
- The control muffins were based on a standard vanilla muffin recipe.
- All ingredients were bought locally, and the taste panel had 30 test subjects.
- Compared to the control muffin, Roselle received the highest overall acceptability score (8.3).
- An average score of 7 or more on a nine-point scale approaches what the researchers call an "extremely acceptable sensory quality."
- Products that achieve such a high score will be close to the quality that food manufacturers want to offer.
- Compared to the control muffin, the protein and fat content of the Roselle muffin was significantly lower. The carbohydrate content was very similar.
- The ascorbic acid content was 12.1 mg/100 g. The recommended daily dose for adults is 75 mg for women and 90 mg for men.

Maybe not so healthy

The Roselle calyx extract obviously provides health benefits, but the muffin research is primarily good news for those of us who already eat muffins. Because there are some bitter tastes — figuratively speaking — in this muffin batter, in the form of unhealthy sugar, saturated fat and white flour.

Eva Falch is an associate professor at NTNU's Department of Biotechnology and Food Science, and a co-author on the study. She doubts that Roselle would be able to be classified as healthy in Norway.



Underused resource

“Roselle calyx extract is an underutilized resource, and the study primarily shows the potential of the plant extract. The researchers use the muffin as a model to say something about how ingredients and steps in the manufacturing process affect and change the properties of the final product,” Falch says.

Eva Falch
is an
associate
professor
and head of
the NTNU
Food
Forum.
Photo:
NTNU

New possibilities

Classifications and health recommendations, as well as taste and food culture, vary between countries.

“In food cultures where baked goods like this are part of the daily diet, Roselle muffins can contribute to increased nutritional value. To make a healthy version, the whole composition should be as good as possible, with little sugar, salt, saturated fat, and so on,” said Falch, who is also head of the NTNU Food Forum.

Animal models and a start-up

A next step could be to test this ingredient in products with a better nutritional profile, such as whole grain products, so that it can be part of a healthy diet. Professor Nutan Kaushik, says the findings have helped in getting the acceptance among consumers as well as regulators.

“Next, we plan to do study the health benefits on animal model and launch a start-up company,” she says.

Moist muffins mould easily

Shelf life was the third factor the researchers investigated. When dry and semi-dry baked goods such as biscuits are left for too long, they can deteriorate both physically and chemically. Muffins with a higher moisture content, on the other hand, are more vulnerable to bacteria, fermentation and mould. Mould and mould spores are destroyed by heat during baking, but as soon as the muffins are out of the oven, they are exposed to airborne mould spores.

... but Roselle keeps fresh without preservatives

Roselle muffins have no added preservatives. The beauty is that the antioxidant ascorbic acid is also found in the pastry. This is a chemical bond that can contribute to increased shelf life in food products. And when it comes to the effects of ascorbic acid

Media Coverage of Roselle Muffin

on the human body, most of us know it by its common name, vitamin C. The tests showed that Roselle kept well for six days at room temperature, with no signs of mould or spores.

Reference: Marak, S.; Kaushik, N.; Dikiy, A.; Shumilina, E.; Falch, E.

Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology Foods 2022, 11, 3982.



Home » Disability » Healthy Roselle Muffins Containing Hibiscus Extract



DISABILITY

Healthy Roselle Muffins Containing Hibiscus Extract



By Editor – March 8, 2023

No Comments

7 Mins Read

Release: 2023-03-07

author: Norwegian University of Science and Technology | contact: ntnu.edu

Peer-reviewed publications: Yes | Doi: <https://www.mdpi.com/2304-8158/11/24/3982>

Additional References: nutrition and health food publications



*Synopsis: The new Supermuffin was named Roselle because it contains calyx extract from the tropical plant *Hibiscus sabdariffa*, which is often called by the same name. , polyphenols, flavonoids, betaine, hibiscus acid, and many other valuable bioactive compounds. It can be used for developing functional foods.*

advertisement

meaning

muffin

Muffins are individually portioned baked goods. However, the term he may refer to either of two specific items. “English-style” flatbreads that are partially raised, such as crumpets, are baked and usually cooked on an unsweetened griddle or “American-style”, often chemically fermented and then sweetened. ‘s quick bread. Baked in a mold. The flatbread

variety is of British or other European derivation and dates back to at least his early 18th century, while the quickbread he originated in North America in the 19th century.

main digest

Enriched muffins from roselle calyx extract using response surface methodology – food.

Health and muffins aren't two words we usually associate with each other, but botanical extracts may help change that at least a little.

do you like muffins? We're talking delicious fluffy muffins that are free of artificial additives and packed with lots of beneficial nutrients. The journal describes the development of just such a new muffin *food*.

From muffins to functional foods

The new supermuffin was named roselle because it contains a calyx extract from the tropical plant hibiscus sabdariffa, which is often called by the same name.

Here, let's take a look at the various formulas tested by researchers.

Hibiscus flowers have a beautiful red color, and calyx extracts contain many valuable bioactive compounds such as polyphenols, flavonoids, betaine, and hibiscus acids. It can be used to develop new products and functional foods that can reduce and potentially have a positive impact on health.

The study also shows that roselle muffins stay fresh for up to six days at room temperature, but with so many beneficial properties, they may be gone long by then.

health benefits

The research is part of Re-FOOD, a joint Norwegian-Indian project focused on improving the use and value of leftover raw materials in food processing.

Professor Nutan Kaushik of Amity University, Uttar Pradesh, India, points out the health benefits of antioxidants, especially in neutralizing free radicals, which can be a key factor in serious illness. I'm here.

prefer natural coloring

Media Coverage of Roselle Muffin

5/1/23, 1:37 PM

Healthy Roselle Muffins Containing Hibiscus Extract – USA Health News

Roselle is also rich in anthocyanins. This dye dissolves in water and can be red, pink, purple, blue, blue, purple, or violet. The color of food greatly influences whether we perceive it as delicious.

Artificial dyes are often clearer and stronger than natural substances. However, concerns about artificial dyes harming our health have increased the demand for natural dyes.

Tasty, nutritious and long shelf life

In the muffin study, 30 people tested 17 recipe variations.

The aim was to find the best roselle muffins based on three key characteristics:

- Muffins should have valuable nutritional properties.
- Muffins should have a taste and texture that people like.
- Muffins should have the best possible shelf life without preservatives.

Aim for the best possible response

Researchers measured and investigated subjects' responses to pastry color, appearance, aroma, texture, fluffiness, elasticity, freshness, and resistance to chewing. We modeled and analyzed how subject responses were influenced by various factors. This method aims to achieve the most positive response to what is tested.

Figure 1. Images of 17 roselle muffins obtained with different formulation combinations. Researchers tested different formulations of muffins containing botanical extracts to see which varieties appealed the most. Screenshot: Nutritionally enriched muffins from roselle calyx extract using response surface methodology. food. 2022; 11(24):3982.

Natural coloring good for cells

Nutritionally, the results suggest that hibiscus extract provides muffins with high levels of antioxidants, such as phenols.

These are substances that are believed to be able to prevent processes in the body that can damage cell membranes and the genetic material within cells. There are thousands of different phenols, most of them found in fruits, juices and wine. , cocoa, tea, coffee, vegetables, flaxseeds, whole grains, and legumes.

may not be very healthy

Roselle calyx extract offers health benefits, but the muffin research is mostly good news for people who already eat muffins. Because, figuratively speaking, it has some bitterness.

Eva Falch is an associate professor in NTNU's School of Biotechnology and Food Sciences and co-author of the study. She wonders if Roselle can be classified as healthy in Norway.

underutilized resources

“Roselle calyx extract is an underutilized resource, and research has largely shown the potential of plant extracts. Researchers used muffins as a model to determine the ingredients and steps in the manufacturing process. It says something about how it affects and changes the properties of the final product,” says Falch.

new possibilities

Classification and health recommendations, as well as taste and food culture, vary from country to country.

“In a food culture where baked goods like this are part of the daily diet, roselle muffins can contribute to the nutritional value of the muffin. To create a healthier version, sugar, salt, saturated fat, etc.” said Falch, who also heads the NTNU Food Forum.

animal models and start-ups

The next step is to test this ingredient in products with better nutritional profiles, such as whole grain products, so that it can be part of a healthier diet.

Professor Nutan Kaushik said the findings have gained acceptance among consumers and regulators.

“We will then study the health benefits of animal models and launch a start-up company,” she says.

Moist muffin Easy to get moldy

Shelf life was the third factor investigated by researchers.

Media Coverage of Roselle Muffin

5/1/23, 1:37 PM

Healthy Roselle Muffins Containing Hibiscus Extract – USA Health News

Dry and semi-dry baked goods such as biscuits can deteriorate physically and chemically if left for too long. Muffins with a high moisture content, on the other hand, are more susceptible to bacteria, fermentation, and mold. exposed to mold spores inside.

Roselle stays fresh without preservatives

Roselle muffins have no added preservatives.

The beauty is that the antioxidant ascorbic acid is also present in the pastry. Most people know the effects of ascorbic acid on the human body by its generic name, vitamin C. Tests showed that Roselle at room temperature she kept well for 6 days with no signs of mold or spores.

reference

Malak, S. Kaushik, N.; Dikiy, A.; Shumilina, E.; Falch, E. Fortified muffins from roselle calyx extract using response surface methodology Foods 2022, 11, 3982.

Source:

Healthy Roselle Muffin with Hibiscus Extract | | Norwegian University of Science and Technology (ntnu.edu). Disabled World makes no warranties or representations in connection therewith. Content may have been edited for style, clarity, or length.

[Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology Foods 2022, 11, 3982.](#)

Bakery&Snacks

By **Oliver Morrison**

07-Mar-2023 - Last updated on 07-Mar-2023 at 15:49 GMT

RELATED TAGS [Muffin](#) [Nutrition](#) [Bakery](#)

Norwegian academics have demonstrated how using the tropical plant *Hibiscus sabdariffa* can contribute to increased nutritional value in everyday foods.



Adaptogens and botanical flavours like hibiscus are on the rise in NPD thanks to their health and wellness credentials. Up to now however, hibiscus has tended to be employed in beverage formulations.

Norwegian researchers have now used the plant extract to unveil a 'super muffin' which contains no artificial additives and a host of beneficial nutrients.

The new 'tasty and fluffy' muffin has been named Roselle, because it contains calyx extract from the tropical plant *Hibiscus sabdariffa*, which is often referred to by the same name.

Hibiscus flowers have a beautiful red colour, and the calyx extract contains many valuable bioactive compounds, such as polyphenols, flavonoids, betaine and hibiscus acid. All of these components can be used to develop new products and functional food that may offer positive health effects, possibly also reducing the risk of chronic diseases, according to the researchers.

Roselle is also rich in anthocyanins. This is a dye that can be dissolved in water and yields a red, pink, purple, blue, blue violet, or violet colour, to add a strong influence on how appetizing people think it is. Artificial dyes are often clearer and more intense than natural substances. But the demand for natural dyes is increasing because of the concern that artificial dyes are harmful to our health.

The study, published in the journal *Food*, also showed that Roselle muffins maintain their freshness for up to six days at room temperature.

The study was part of Re-FOOD, a Norwegian-Indian collaborative project focusing on using and enhancing the value of rest raw materials of food processing.

In the muffin study, 30 people tested a total of 17 recipe variants. The aim was to find the best version of Roselle muffins based on three important qualities: taste and texture; valuable nutritional properties; and the best possible shelf life without preservatives.

Bakery&Snacks

The hibiscus extract gave muffins high levels of antioxidants

In terms of nutrition, the results suggested that the hibiscus extract provides muffins with high levels of antioxidants like phenolics. These are substances believed to be able to prevent processes in the body that can damage cell membranes or the genetic material in our cells. There are several thousand different phenols, most of which are found in fruit, juice, wine, cocoa, tea, coffee, vegetables, flax seed, wholegrains and legumes.

While the Roselle calyx extract gave health benefits to muffins, the study didn't produce what could be described a 'healthy muffin', stressed Eva Falch, an associate professor at NTNU's Department of Biotechnology and Food Science. That's because the muffin batter still used unhealthy sugar, saturated fat and white flour. "Roselle calyx extract is an underutilized resource, and the study primarily shows the potential of the plant extract. The researchers use the muffin as a model to say something about how ingredients and steps in the manufacturing process affect and change the properties of the final product," Falch said.

"In food cultures where baked goods like this are part of the daily diet, Roselle muffins can contribute to increased nutritional value. To make a healthy version, the whole composition should be as good as possible, with little sugar, salt, saturated fat, and so on," she added.

Moist muffins mould easily but Roselle keeps fresh without preservatives

Shelf life was the third factor the researchers investigated. When dry and semi-dry baked goods such as biscuits are left for too long, they can deteriorate both physically and chemically. Muffins with a higher moisture content, on the other hand, are more vulnerable to bacteria, fermentation and mould. Mould and mould spores are destroyed by heat during baking, but as soon as the muffins are out of the oven, they are exposed to airborne mould spores.

The Roselle muffins have no added preservatives. According to the researchers, the beauty is that the antioxidant ascorbic acid is also found in the pastry. This is a chemical bond that can contribute to increased shelf life in food products. And when it comes to the effects of ascorbic acid on the human body, most of us know it by its common name, vitamin C. The tests showed that Roselle kept well for six days at room temperature, with no signs of mould or spores.

Reference

Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology

Published Journal - Foods



Article

Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology

Sengnolotha Marak ^{1,2}, Nutan Kaushik ^{1,*}, Alexander Dikiy ³, Elena Shumilina ³ and Eva Falch ³

¹ Amity Food and Agriculture Foundation, Amity University Uttar Pradesh, Noida 201313, UP, India

² Amity Institute of Food Technology, Amity University Uttar Pradesh, Noida 201313, UP, India

³ Department of Biotechnology and Food Sciences, Norwegian University of Science and Technology (NTNU), 7491 Trondheim, Norway

* Correspondence: nkaushik5@amity.edu.in; Tel.: +91-98-1139-2249

Abstract: *Hibiscus sabdariffa*, often called Roselle, is a flowering plant with a variety of traditional medicinal uses. Its calyx, with a bright and attractive red color, produces a tart and pleasant acidic taste. The purpose of this study was to develop a Roselle muffin and assess the acceptability, nutrition, and shelf life of the muffin using its ingredients. The muffin was developed using different formulations in different proportions resulting from Response Surface Methodology (RSM). Sensory parameters were used to assess the muffin's acceptability. According to the findings, the combination of extract volume 45.37 mL, citric acid 1.11 g, and sodium bicarbonate 1.67 g produces the best muffin, with the panelist's sensory scores reaching up to 84%. The outcome of the study suggests muffins baked with the Roselle calyx extract have high antioxidant ($12.53 \pm 0.13\%$), anthocyanin (126.63 ± 1.96) mg Cyn-3-glu/100 g, phenolic (12.91 ± 0.69) mg GAE/100 g, and ascorbic acid (12.10 ± 0.89) mg/100 g contents. The microbial shelf life of the developed muffin is estimated to be 6 days at room temperature. The study findings can therefore be utilized in the development of foods containing Roselle calyx extract.

Keywords: *Hibiscus sabdariffa*; Roselle calyx extract; Roselle muffin; response surface methodology



Citation: Marak, S.; Kaushik, N.; Dikiy, A.; Shumilina, E.; Falch, E. Nutritionally Enriched Muffins from Roselle Calyx Extract Using Response Surface Methodology. *Foods* **2022**, *11*, 3982. <https://doi.org/10.3390/foods11243982>

Academic Editor: Francisca Rodrigues

Received: 3 November 2022

Accepted: 6 December 2022

Published: 8 December 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Due to their bioactive components, adding plant extracts to food products has become more popular. Natural bioactive components have a significant impact on daily activities and are linked to numerous health advantages and minimal toxicity. Many diseases have been treated with plant-derived medicines because of their long-recognized therapeutic properties [1]. In the development of functional foods and the treatment of human diseases, several active substances with various biological effects are widely used [2–4]. An excellent source of molecules for the development of nutraceuticals, functional foods, and food additives are offered by natural bioactive compounds with a wide variety of structural and functional properties [5].

The aqueous extract of *H. sabdariffa* calyces contains a wide spectrum of several groups of bioactive compounds, including anthocyanins, polysaccharides, flavonoids, and other organic compounds including betaine and hibiscus acid [6]. Anthocyanins, which are abundant in calyces and are responsible for their red pigmentation, have undergone extensive scientific research owing to their distinctive color characteristics [7,8]. Numerous studies have demonstrated that *H. sabdariffa* and its extract exhibit useful qualities and water-soluble bioactive components that can be exploited to develop new products with additional nutritional attributes that might enhance consumers' health [6,9,10]. Consumer acceptability of any dish is greatly influenced by its color. Natural colors have become more prominent due to a growing public concern over the toxicity of synthetic colorants [11]. The calyces of *H. sabdariffa* are thought to be excellent and attractive sources of red colorants that are water-soluble and could be used as natural food colorings [12,13]. In parts of the

world including West Africa and South Asia, *H. sabdariffa* is used to make tea. It has a tart flavor, and it is sold commercially in these regions as jams and juices [7].

While developing any bakery products, quality parameters such as texture, phytochemical content, and shelf life are important to consider. Muffins that are well-aerated, elastic, and possess a degree of springiness are high-quality and fresh [14]. Another important texture parameter is chewiness, which is mostly related to how tough it is to chew before swallowing [15].

In addition to offering basic nutrition, phytochemicals present in fruits, vegetables, grains, and other plant foods may also provide health advantages by lowering the risk of serious chronic diseases [16]. However, during processing, these phytochemicals are subjected to a variety of degradations. The physical characteristics and chemical composition of foods change because of cooking, which makes this alteration quite evident [17]. Due to its processing conditions, the results on the changes in phytochemicals during household cooking were inconsistent and occasionally conflicting. For example, Blessington et al. found that cooking potatoes in various ways—including boiling, baking, frying, and microwaving—significantly raises their overall phenolic content and antioxidant activity [18]. On the contrary, Xu et al. discovered that boiling, baking, and microwaving decreased phytochemical concentrations and their antioxidant activity [19].

Shelf life is a particularly important aspect in the production of food products. Low- and medium-moisture bakery items are more prone to physical and chemical deterioration, whereas high-moisture products are more vulnerable to microbiological decomposition by bacteria, yeast, and molds. Although the heat while baking destroys mold and mold spores by thermal reactivity, post-baking contamination from airborne mold spores occurs during handling procedures such as cooling and packing [20]. According to Gilbert et al., cakes with aerobic plate counts greater than 6 log₁₀ CFU had unacceptable microbiological quality, whereas acceptable values ranged from 5 log₁₀ CFU to 6 log₁₀ CFU [21]. The generic microbiological standard for cakes and pastries, which was published by the Institute of Food Science and Technology (IFST), specified that the maximum permissible levels of yeast and molds are 5 log₁₀ CFU and 4 log₁₀ CFU, respectively [22].

A combination of mathematical and statistical methods, such as the Response Surface Methodology (RSM), is a helpful tool for developing, improving, and optimizing processes [23–26]. This technique has been employed by numerous studies to improve bakery goods [27–29]. Although RSM has been adopted in numerous research as an optimization tool for bakery goods, the optimization of Roselle muffins has not been studied. Therefore, the study's objective was to formulate and optimize the ingredient levels of the Roselle muffin using RSM. Additionally, the phytochemical analysis of the batter and muffins was investigated to determine their retention after baking.

2. Results

2.1. Effect of Factors on Muffin Responses

In this study, the effect of Roselle calyx extract, citric acid, and sodium bicarbonate on sensory responses was observed using the Response Surface Methodology (RSM). The average score of 30 responders and the effect of varying factors (extract volume, citric acid, and sodium bicarbonate concentration) on responses (texture and overall acceptability) is presented in Table 1. The texture score ranged from 6.36 to 7.23, while the overall acceptability (OAA) score was between 5.70 and 7.80. Table S1 provides results for other sensory aspects, such as appearance, color, and taste. Figure 1 shows the color of the Roselle muffin when various formulations are combined.

Following up on these observations, analysis of variance (ANOVA) was performed (Table 2). The quadratic model, with an R² value of 0.70 for texture and 0.94 for OAA, was suggested.

Table 1. Processed data on the muffin score given by the respondents.

Codes	Independent Variables			Dependent Variables			
	A	B	C	Response 1: Texture (Score)		Response 2: OAA (Score)	
Formulation	Extract Volume (mL)	Citric Acid (g)	Sodium Bicarbonate (g)	Experimental Data	RSM Predicted Data	Experimental Data	RSM Predicted Data
01	27.5	2.5	1.0	6.36 ± 1.54	6.54	5.70 ± 1.78	5.88
02	5.0	1.0	1.7	6.63 ± 0.92	6.90	6.00 ± 1.11	6.14
03	27.5	1.7	1.7	6.93 ± 1.11	6.96	7.53 ± 0.97	7.56
04	50.0	2.5	1.7	6.60 ± 1.28	6.47	6.65 ± 1.25	6.50
05	27.5	1.0	1.0	7.23 ± 0.97	6.96	6.53 ± 1.27	6.35
06	5.0	1.7	2.5	6.46 ± 1.00	6.51	6.26 ± 0.94	6.30
07	50.0	1.7	2.5	6.63 ± 1.35	6.58	6.78 ± 0.99	6.99
08	27.5	2.5	2.5	7.13 ± 1.27	7.00	6.66 ± 1.24	6.59
09	50.0	1.7	1.0	6.63 ± 1.09	6.76	6.70 ± 1.08	6.66
10	27.5	1.7	1.7	6.88 ± 0.63	6.96	7.30 ± 0.83	7.56
11	27.5	1.7	1.7	7.23 ± 0.72	7.05	7.80 ± 0.80	7.56
12	27.5	1.7	1.7	7.00 ± 0.69	6.96	7.60 ± 0.77	7.56
13	50.0	1.0	1.7	7.20 ± 1.03	7.12	7.10 ± 0.99	7.07
14	5.0	1.7	1.0	6.96 ± 1.15	6.69	6.38 ± 1.32	6.17
15	27.5	1.0	2.5	7.23 ± 0.85	7.36	6.53 ± 0.93	6.60
16	5.0	2.5	1.7	6.46 ± 1.16	6.54	6.20 ± 1.12	6.23
17	27.5	1.7	1.7	6.76 ± 0.67	6.96	7.56 ± 0.62	7.56



Figure 1. Images of Roselle muffin obtained with different formulations combinations.

This quadratic relationship was observed between the sensory responses and the extract volume, citric acid, and sodium bicarbonate.

$$Y = 6.96 + 0.06 A - 0.21 B + 0.03 C - 0.10 AB + 0.12 AC + 0.19 BC - 0.27 A^2 + 0.04 B^2 - 0.01 C^2 \quad (1)$$

$$Y = 7.56 + 0.29 A - 0.11 B + 0.11 C - 0.16 AB + 0.05 AC + 0.24 BC - 0.44 A^2 - 0.62 B^2 - 0.01 C^2 \quad (2)$$

Equation (1) shows the response texture of the optimal quadratic polynomial equation obtained from RSM. Where Y = texture (score), A, B, and C are extract volume, citric acid, and sodium bicarbonate, respectively.

Equation (2) shows the response OAA of the optimal quadratic polynomial equation obtained from RSM. Where $Y = \text{OAA (score)}$, A, B, and C are extract volume, citric acid, and sodium bicarbonate, respectively.

Furthermore, the model was determined to be significant for OAA (0.01) and not significant for texture (0.21) using $p < 0.05$ standards. This indicates that the panelists' preference for OAA was significantly influenced by variations in the Roselle muffin formulation. OAA (0.22) and texture (0.13) were found to have a non-significant Lack of Fit ($p < 0.05$). Non-significant Lack of Fit is favorable because it implies that the response data and the model are compatible [30].

The diagnostic plots of the experiment and model results show the accuracy of the models for Texture (a) and OAA (b) responses (Figure S1).

Given that the points are rationally close to the straight line and no variance was seen, the normal % probability of residual plot for response is normally distributed in both models.

Predicted values are close to the experimental values which shows a good relationship between the actual and the predicted values. In Figure S1b, the lowest OAA response value is 5.7 which is indicated by blue; meanwhile, the highest OAA response value is 7.8 which is indicated by red color. This can be interpreted that the actual results would be close to the results predicted by the Design Expert program.

Table 2. Result of analysis of variance (ANOVA) texture response (A) and OAA response (B).

A						B					
Source	Sum of Squares	Degree of Freedom	Mean Square	F-Value	p-Value	Source	Sum of Squares	Degree of Freedom	Mean Square	F-Value	p-Value
Model	1.01	9	0.1125	1.86	0.21 **	Model	5.62	9	0.6244	12.82	<0.01 *
A-Extract Volume	0.0378	1	0.0378	0.62	0.45	A-Extract Volume	0.7140	1	0.7140	14.66	<0.01
B-Citric Acid	0.3784	1	0.3784	6.26	0.04	B-Citric Acid	0.1128	1	0.1128	2.32	0.17
C-Sodium Bicarbonate	0.0091	1	0.0091	0.15	0.70	C-Sodium Bicarbonate	0.1058	1	0.1058	2.17	0.18
AB	0.0462	1	0.0462	0.76	0.41	AB	0.1056	1	0.1056	2.17	0.18
AC	0.0625	1	0.0625	1.03	0.34	AC	0.0100	1	0.0100	0.20	0.66
BC	0.1482	1	0.1482	2.45	0.16	BC	0.2304	1	0.2304	4.73	0.06
A ²	0.3242	1	0.3242	5.37	0.05	A ²	0.8441	1	0.8441	17.33	<0.01
B ²	0.0067	1	0.0067	0.11	0.74	B ²	1.63	1	1.63	33.52	<0.01
C ²	0.0007	1	0.0007	0.01	0.91	C ²	1.42	1	1.42	29.10	<0.01
Residual	0.4230	7	0.0604			Residual	0.3410	7	0.0487		
Lack of Fit	0.3012	3	0.1004	3.30	0.13	Lack of Fit	0.2133	3	0.0711	2.23	0.22
Pure Error	0.1218	4	0.0305			Pure Error	0.1277	4	0.0319		
Cor Total	1.44	16				Cor Total	5.96	16			
Fit Statistics	Std. Dev.	Mean	C.V%	R ²		Fit Statistics	Std. Dev.	Mean	C.V%	R ²	
	0.2458	6.84	3.59	0.70			0.2207	6.78	3.25	0.94	

OAA: Overall acceptability analysis; * significant; ** non-significant.

2.2. Optimization

Optimization was conducted to obtain responses that are consistent with the responses that the panelists could be perceived. The components and goals in the optimization stage are shown in Table 3. Maximizing the extract volume, texture, and OAA within a range of citric acid and sodium bicarbonate components was the aim of the optimization process.

Table 3. Components and optimization goal.

Components	Goal	Lower Limit	Upper Limit
A: Extract Volume	Maximize	5.00	50.0
B: Citric Acid	is in range	1.00	2.50
C: Sodium Bicarbonate	is in range	1.00	2.50
Texture	Maximize	6.36	7.23
Overall Acceptability	Maximize	5.70	7.80

According to the Design Expert program's optimization procedure, the following contents were revealed: Citric acid (1.11 g), sodium bicarbonate (1.16 g) and 45.37 mL of the extract volume were present, and the desirability was 0.84 (Table 4). It produces a response score of 8.31 for texture and 8.30 for OAA under such conditions, which is higher than the value predicted by the model (7.13 for texture and 7.27 for OAA). Therefore, it can be stated that optimization and standardization using RSM were successful in the development of the Roselle muffin.

Table 4. The optimization and outcome of the optimized responses.

Components	Optimum Value	
Extract Volume (mL)	45.37	
Citric acid (g)	1.11	
Sodium bicarbonate (g)	1.67	
Desirability of response value	0.84	
Response	Model Calculated value	Observed value
Texture	7.13	8.31
OAA	7.27	8.30

2.3. Sensory and Physicochemical Characteristics

The sensory evaluation scores for color and appearance, aroma, body and texture, taste and flavor, and overall acceptability that were received from the semi-trained panelists are presented in Table 5. The muffin's sensory attributes were valued. Some panelists stated that the Roselle muffin tasted slightly sour, setting it apart from other muffins sold commercially. The high concentration of organic acids, particularly hibiscus acid in Roselle calyx, is what gives muffins their sour flavor [6]. In comparison to the control muffin, the Roselle muffin had higher acceptability scores for attributes, particularly in body and texture (8.21) and taste and flavor (8.18). Similar findings were reported in the study of Siti Faridah et al., which determined that the panelists regarded quality scores with 10% Roselle calyx powder to be the most acceptable [31]. When compared to the control muffin in our study, the muffin enhanced with Roselle calyx extract (not powder) received the highest overall acceptability score (8.3). The addition of Roselle extract enhanced the acceptability. In sensory evaluation, an average liking score of 7 or more on a nine-point scale often indicates extremely acceptable sensory quality [32]. Hence, a product obtaining this score may be taken into consideration as a competent description of the desired quality.

The physicochemical parameters of muffins are presented in Table 5. Roselle calyx extract did not significantly ($p < 0.05$) increase the moisture content of the Roselle muffin. The ash concentration in the Roselle muffin was found to be statistically ($p < 0.05$) higher than that of the control muffin. The high content of ash in the Roselle muffin may be due to the rich amount of calcium, iron, and crude fiber in Roselle calyx. However, compared to the control muffin, the protein and fat content of the Roselle muffin revealed a considerably ($p < 0.05$) lower value. The total carbohydrate content between the Roselle muffin and the control muffin is not significantly different ($p < 0.05$). Comparing the Roselle muffin to the control muffin, the pH of the Roselle muffin is slightly acidic (5.26). Ascorbic acid is present in reasonable amounts in the Roselle muffins (12.1 mg/100 g). For adult women and men, the Dietary Recommended Allowance (RDA) of ascorbic acid is 75 mg per day and 90 mg per day, respectively, to offer antioxidant protection [33].

Color is an important parameter affecting the overall acceptability of the product. Color analysis shows that a Roselle muffin had a reddish color with L^* (62.70 ± 0.19), a^* value (9.51 ± 0.01) and b^* value (10.47 ± 0.06), and that of the control muffin was a brownish-yellow color with L^* (68.84 ± 0.21), a^* value (5.21 ± 0.05), and b^* value (14.88 ± 0.08). In this study, we see that the Roselle muffin had a lighter (L^*) color compared to the control muffin with higher a^* values indicating a red, dark color with a pleasant taste due to the presence of the Roselle calyx extract.

The texture parameters as revealed by instrumental analysis are summarized in Table 5. In every parameter studied, the textural profile of the Roselle muffin showed a significant variation. The Roselle muffin's texture was significantly changed by sodium bicarbonate and citric acid, giving it a better texture profile than the control muffin. The hardness of the Roselle muffin was reduced by the addition of citric acid and sodium bicarbonate from 1720.84 N in the control to 865.38 N in the Roselle muffin. The volume of the crumb and the overall volume of air cells are connected to hardness [34]. With rising gas cell size, the crumb structure of the product becomes softer. Springiness was observed in the control muffin at 0.89% and the Roselle muffin at 0.87%. There were no obvious changes in the springiness of the control and Roselle muffins. Cohesiveness, which describes how well a food retains its form between the first and second chew, showed a significant difference between the control and Roselle muffins. With the addition of citric acid and sodium bicarbonate, the value of chewiness in the Roselle muffin significantly decreased.

Table 5. Sensory and physicochemical characteristics of muffin.

Parameters	Roselle Muffin	Control Muffin	p-Value ($p < 0.05$)
Sensory Parameters			
Colour and Appearance	7.85 ± 0.70 *	7.36 ± 0.80 *	0.01
Aroma	7.71 ± 0.66 ^{ns}	7.70 ± 0.70 ^{ns}	0.92
Body and Texture	8.21 ± 0.53 **	7.53 ± 0.89 **	<<0.01
Taste and Flavour	8.18 ± 0.70 *	7.76 ± 0.81 *	0.03
Overall Acceptability	8.30 ± 0.68 **	7.68 ± 0.64 **	<<0.01
Proximate (per 100 g)			
Moisture (%)	21.66 ± 0.89 *	20.86 ± 0.24 *	0.29
Ash (%)	4.22 ± 0.41 **	2.11 ± 0.41 **	<<0.01
Fat (%)	11.20 ± 0.16 **	13.6 ± 0.16 **	<<0.01
Protein (%)	7.73 ± 0.12 **	8.33 ± 0.12 **	<<0.01
Total Carbohydrate (%)	55.18 ± 1.24 ^{ns}	55.09 ± 0.51 ^{ns}	0.92
pH	5.26 ± 0.04 **	7.10 ± 0.08 **	<<0.01
Ascorbic acid, mg/100 g	12.10 ± 0.89	ND	-
Color Parameters			
<i>L</i> *	62.70 ± 0.19 **	68.84 ± 0.21 **	<<0.01
<i>a</i> *	9.51 ± 0.01 **	5.21 ± 0.05 **	<<0.01
<i>b</i> *	10.47 ± 0.06 **	14.88 ± 0.08 **	<<0.01
<i>c</i> *	14.14 ± 0.05 **	15.77 ± 0.10 **	<<0.01
<i>h</i> ⁰	47.75 ± 0.11 **	70.69 ± 0.06 **	<<0.01
Texture Parameters			
Hardness, N	865.38 ± 6.32 **	1720.84 ± 56.08 **	<<0.01
Springiness, %	0.87 ± 0.01 ^{ns}	0.89 ± 0.00 ^{ns}	0.30
Cohesiveness	0.45 ± 0.00 *	0.50 ± 0.01 *	0.02
Chewiness	346.26 ± 1.71 **	780.11 ± 41.87 **	<<0.01

ND = not detected; * significant; ** highly significant; ^{ns} non-significant.

2.4. Phytochemical Characteristics of Batter and Optimized Muffin

The phytochemical attributes of batters and Roselle muffins are presented in Table 6. In total, the batter contained 154.56 mg C3G of anthocyanins (TAC / 100 g). The TAC was 126.63 mg Cyn-3-glu/100 g after baking; which suggests that about 82% of the TAC was retained. The total phenolic content (TPC) in the batter for the Roselle muffin significantly decreased from 19.74 to 12.91 mg GAE/100 g. Indicating that only 65% of the TPC was retained. The antioxidant activity of the batter was 27.30%, while that of the Roselle muffin was 12.53%. Thus, the reduction in antioxidant activity was nearly 50%. There was a modest reduction in TAC and antioxidant properties after baking.

Table 6. Phytochemical content of batter and muffin.

Phytochemical Parameters	Roselle Muffin		Control Muffin		p-Value (<i>p</i> < 0.05)
	Batter	Muffin	Batter	Muffin	
TAC, mg cyanidin-3-glucoside (Cyn-3-glu/100 g)	154.56 ± 3.40 **	126.63 ± 1.96 **	ND	ND	<<0.01
TPC, mg gallic acid (GA/100 g)	19.74 ± 0.28 **	12.91 ± 0.69 **	3.12 ± 0.06	ND	<<0.01
Antioxidant activity (%)	27.30 ± 1.03 **	12.53 ± 0.13 **	ND	ND	<<0.01

TAC: Total anthocyanin content; TPC: Total phenolic content; GA: Gallic acid; ND: Not Detected; ** highly significant.

2.5. Microbial Shelf-Life

In this study, the muffins’ microbiological shelf-life was examined every three days and are shown in Table 7. In the first and third days of room-temperature storage (25 °C), the overall plate counts were below the limit for detection. The sixth day of storage saw the appearance of a few bacterial colonies in the Roselle muffin. After the sixth day of storage, there were an infinite number of colonies. The analysis was stopped on day 12 based on the observation of microbial growth, which showed that molds had begun to form on the ninth day of storage in the Roselle muffin. According to this study, during the ninth day of storage at room temperature, both the total plate count and the yeast and mold count increased in the Roselle muffin. However, the control muffin is within the permissible limit up to day 9 of the storage. This may be due to the presence of anthocyanins in Roselle calyx extract. Microencapsulation may be the solution for improving the stability of anthocyanins [35]. Aerobic plate counts acceptable values ranged from 5 log₁₀ CFU to 6 log₁₀ CFU [21] and maximum permissible levels of yeast and molds 5 log₁₀ CFU 4 log₁₀ CFU, respectively [22]. Because the results of the microbiological count were higher than the permissible limit, the findings of the present study suggest that Roselle muffins may not be safe to consume after the sixth day of storage at room temperature. It is mostly due to no preservatives added to the muffin. Without preservatives, these microorganisms will eventually degrade baking products, reducing their shelf life. Our study is in line with that of Rodriguez et al., who discovered that cake samples with no preservative were moldy on the surface after 6 days of storage at 15–20 °C [36].

Table 7. Total plate count and yeast and molds during storage of muffins (cfu/g).

Days	Roselle Muffin		Control Muffin	
	Total Plate Count	Yeast and Molds	Total Plate Count	Yeast and Molds
01	2.4 × 10 ³	4.1 × 10 ³	3 × 10 ²	2 × 10 ²
03	1.8 × 10 ⁴	1.5 × 10 ⁴	3 × 10 ²	5 × 10 ²
06	5.4 × 10 ⁴	4.9 × 10 ⁴	7 × 10 ³	3 × 10 ³
09	Uncountable	Uncountable	5.7 × 10 ⁴	4.5 × 10 ⁴
12	Visible mold growth	Visible mold growth	Uncountable	Uncountable
15	Visible mold growth	Visible mold growth	Visible mold growth	Visible mold growth

3. Materials and Methods

3.1. Chemicals

Trichloroacetic acid (99%), Gallic acid (98%), and 2,2-diphenyl-1-picrylhydrazyl (DPPH, 95%) were obtained from Sisco Research Laboratory Pvt. Ltd. (Mumbai, India). L-ascorbic acid (99%); sodium carbonate, anhydrous (99.5%); glacial acetic acid (99.5%); and petroleum ether were purchased from Thermo Fisher Scientific (Mumbai, India). 2,6-dichlorophenol-indophenol sodium salt (DCIP) and the Folin & Ciocalteus Phenol (FCP) reagent were obtained from Central Drug House (P) Limited (New Delhi, India);

nutrient agar and potato dextrose agar (PDA) were obtained from HiMedia; methanol and ethanol of analytical grade were procured from Rankhem (New Delhi, India).

3.2. Raw Materials

The research material was centered on the development of nutritionally enriched muffins. In the current study, dried Roselle (*H. sabdariffa*) calyces from the Garo Hills in Meghalaya, India, were used as a source of natural pigment [6]. Flour, sugar, butter, eggs, salt, citric acid powder, and sodium bicarbonate have also been acquired at the neighborhood store in order to carry out the muffin-making process.

3.3. Design of the Experiment

The experiment was planned using the Response Surface Methodology (RSM) Box–Behnken’s Design (BBD). Operationally, the method was designed through data processing by applying the Design Expert (DX) Program Version 13.0.6, Stat-Ease Inc., Minneapolis, MN, USA (www.statease.com (accessed on 23 August 2021)). Three independent factors (variables), namely, extract volume, citric acid, and sodium bicarbonate, were employed in this study; their respective codes were A, B, and C. The findings of the preliminary study were used to determine the independent variables. The independent variables’ minimum and maximum values were entered into the model for randomization. Table 8 displays every level of the variable included in the model. For the three independent variables, a total of 17 runs were generated after randomization (Table 9).

Table 8. Variable levels for preparing muffins.

Codes	Independent Variables	Units	Minimum	Maximum
A	Extract Volume	mL	5.00	50.00
B	Citric Acid	g	1.0000	2.50
C	Sodium Bicarbonate	g	1.0000	2.50

Table 9. Experimental design for preparation of Roselle muffins based on RSM along with control.

Formulation	Independent Variables				Dependent Variables					
	A: Extract Volume (mL)	B: Citric Acid (g)	C: Sodium Bicarbonate (g)	Milk Volume (mL)	Flour (g)	Sugar (g)	Butter (g)	Egg * (g)	Vanilla Extract (g)	Salt (g)
1	27.5	2.5	1.0	0	100	60	50	100	1	1
2	5.0	1.0	1.7	0	100	60	50	100	1	1
3	27.5	1.7	1.7	0	100	60	50	100	1	1
4	50.0	2.5	1.7	0	100	60	50	100	1	1
5	27.5	1.0	1.0	0	100	60	50	100	1	1
6	5.0	1.7	2.5	0	100	60	50	100	1	1
7	50.0	1.7	2.5	0	100	60	50	100	1	1
8	27.5	2.5	2.5	0	100	60	50	100	1	1
9	50.0	1.7	1.0	0	100	60	50	100	1	1
10	27.5	1.7	1.7	0	100	60	50	100	1	1
11	27.5	1.7	1.7	0	100	60	50	100	1	1
12	27.5	1.7	1.7	0	100	60	50	100	1	1
13	50.0	1.0	1.7	0	100	60	50	100	1	1
14	5.0	1.7	1.0	0	100	60	50	100	1	1
15	27.5	1.0	2.5	0	100	60	50	100	1	1
16	5.0	2.5	1.7	0	100	60	50	100	1	1
17	27.5	1.7	1.7	0	100	60	50	100	1	1
Control	0	0	0	50	100	60	50	100	1	1

* Whole raw beaten egg at room temperature.

3.4. Roselle Calyx Extract Preparation

For extraction, the fine calyx powder (10 g) was heated for 5 min at 80° C after soaking overnight at room temperature in 100 mL of distilled water. The extract was filtered

through filter paper and made the final volume to 100 mL. It was then added to the batter of the muffins.

3.5. Muffin Preparation

The control muffin was based on a simple vanilla muffin recipe [37] with a slight modification (Table 9) while in the treatments, the baking powder was replaced with citric acid and sodium bicarbonate to maintain the pH of the batter as the color of anthocyanin is pH dependent, and the Roselle extract was added in place of milk in the proportion mentioned in Table 9. It involved combining all the specified ingredients in an electric mixer and blending them at high speed for five minutes. The batter was combined, then poured into a muffin pan, and baked for 20 min at 180 °C. (OTG Wonder Chef oven, Wonder Chef, Mumbai, India). Five batches were prepared for the optimized formulation. The standard vanilla muffin recipe was used to make the control muffin. The muffins were packed in a clean metallic polyethylene zip lock bag (32-micron) after baking and cooling in preparation for further testing.

3.6. Optimization

The Design Expert Program's optimization objective was used to evaluate each response (texture and overall acceptability). According to the data from the fitted variables and the response score, optimization was carried out. The Design Expert Program interprets the results of the optimization and generate new optimum formulation.

3.7. Sensory Analysis

The muffins' sensory preferences were assessed using a 9-point hedonic scale, with 1 being highly dislike and 9 representing a favorable opinion (like very much). The samples, which were distributed at random, were analyzed by a semi-trained panel comprising 30 panelists. All the panelists were familiar with food tasting and capable of differentiating the taste, flavor, texture, etc., and presenting their reactions. The panelists ranged in age from 15 to 60. The presentation of the samples was carried out in a random order, and the sample sets were allotted with different codes. Each time on the day of analysis, the panelists were given four sets of different formulations and one control sample for sensory evaluation. Each formulation was tested by 30 panelists ($n = 30$). Taste-neutral water was provided for rinsing, and the samples were presented at room temperature.

3.8. Physicochemical Analysis

For all analyses, three replicates were taken and each replicate consisted of one muffin. The details of the methodology are given below.

3.8.1. Proximate

The proximate composition of the optimized and control muffins in terms of moisture, fat, protein, ash, and total carbohydrates were determined using standardized and validated laboratory AOAC methods [38].

3.8.2. Texture Profile Analysis

The method provided by Jauraha et al. served as the basis for the instrumental analysis of muffin texture [39]. Using the TA-XT-plus Texture Analyzer, a texture profile analysis of the optimized and control muffins was performed at IARI, PUSA, New Delhi, India. For compression, a 75 mm diameter P/35" aluminum plate was employed. The following conditions were used to conduct the test: Test speed was 1 mm/s, the trigger force was 5 g, and the strain was 50%. The muffin was compressed twice to assess its textural characteristics, such as its hardness, springiness, cohesiveness, and chewiness.

3.8.3. PH Measurement

A pH meter (LMPH-10, LABMAN Scientific instrument, Chennai, India) was used to measure the muffin's pH [40]. To separate the solids and liquid, 0.5 g of the sample was treated with 20 mL of distilled water, vortexed for 3 min, and then, left at room temperature for 1 h. The pH of the supernatant was determined after centrifugation for 3 min at 3050× g.

3.8.4. Color Measurement

Using a CIE color measuring instrument (NR100 Precision colorimeter, 3nh, Shenzhen, China) and the $L^* a^* b^*$ color scale system, the measurement of color was analyzed [40]. The a^* value stands for redness/greenness, b^* value for yellowness/blueness, and L^* value for lightness/darkness. For color analysis, 1 muffin was powdered, of which 20 g of a muffin was pulverized into small particles prior to the examination and placed on a particular plate for analysis. The muffin's color was automatically detected and displayed on the screen.

3.9. Phytochemical Analysis

3.9.1. Total Phenolic Content

The Folin–Ciocalteu method was used to determine the total phenolic content (TPC) [41]. In brief, a stock solution of gallic acid (1 mg/mL) was prepared at different concentrations in methanol to produce a standard gallic acid curve. These concentrations were mixed with 2 mL of 10% Folin–Ciocalteu reagent. Then, 1 mL of 10% sodium carbonate solution was added after 6 min. After 90 min, the absorbance was measured at 760 nm using UV–VIS spectrophotometer (LMSP-UV1900, LABMAN). For the sample extracts, the same process was repeated. Gallic acid equivalents (mg GAE/g) were used to calculate the total amount of phenol in the sample.

3.9.2. Total Anthocyanin Content

According to the method provided by Lee et al., the total anthocyanin content (TAC) was determined [42]. A total of 10 mL of distilled water were used to dilute 1 mL of the sample extract. Then, 1 mL of the diluted sample solution was diluted with 5 mL with buffer pH 1.0 into a test tube and wrapped with aluminum foil. Next, 1 mL of the diluted sample solution was further diluted with 5 mL with buffer pH 4.5 into test tube and wrapped with aluminum foil. The absorbance at 520 nm and 700 nm was measured using a UV–VIS spectrophotometer (LMSP-UV1900, LABMAN) in a 4 mL spectrophotometer glass cell after the mixtures were let to stand for 30 min at room temperature. According to the following equation, results were reported as Cyanidin-3-glucoside equivalents per 100 g of sample.

$$CA = (A \times MW \times DF \times 100) / (\epsilon \times 1)$$

CA: Anthocyanin concentration (mg/100 g); A: absorbance difference A (A520 nm–A700 nm) pH 1.0 – (A520 nm–A700 nm) (A520 nm–A700 nm) pH 4.5; MW (Molecular Weight): cyanidin-3-glucoside weighs 449.2 g/mol; DF: dilution factor; 1: pathlength in cm; ϵ : cyanidin-3-glucoside's molar extinction coefficient (26,900); 100: conversion factor for deriving mg/100 g.

3.9.3. DPPH Radical Scavenging

The procedure described by Brand-Williams et al. [43] with a slight modification was used to carry out the DPPH radical scavenging activity. In brief, 5.92 mg of DPPH was dissolved in 100 mL of methanol to make a DPPH solution. Then, 1 mL sample extract and 3 mL of DPPH were mixed, vortexed, and kept in the dark at room temperature for 30 min. The control sample, without any extract, was prepared in the same volume. Then, using methanol as a blank, the absorbance was spectrophotometrically measured at 515 nm using

UV–VIS spectrophotometer (LMP-UV1900, LABMAN). The following equation was used to determine the radical scavenging activity

$$\% \text{ inhibition} = (\text{Abs of control} - \text{Abs of sample} / \text{Abs of control}) \times 100$$

3.9.4. Ascorbic Acid Content

Ascorbic acid content was determined using 2,6-dichlorophenolindophenol (DCIP) dye-titrimetric method [40]. The TCA solution was prepared by dissolving 15 g of trichloroacetic acid (TCA) in 200 mL of distilled water and 40 mL of acetic acid. Thereafter, it was diluted with distilled water to 500 mL and filtered using filter paper. To make a standard ascorbic acid solution, weigh 0.05 g ascorbic acid and dissolve it in 60 mL TCA solution and make a final volume of 250 mL with distilled water. The dye standard solution was prepared by dissolving 0.05 g of 2,6-dichlorophenolindophenol and 0.05 g sodium carbonate (Na_2CO_3) in 100 mL distilled water. The dye was standardized by titrating it against 10 mL of ascorbic acid stock solution until a faint pink color that lasted for a few seconds was obtained

In order to prepare the sample, 0.5 g was weighed and 20 mL of TCA solution was added to it. The mixture was diluted to the desired strength with distilled water and filtered into a volumetric flask. An aliquot of 10 mL was titrated against the reference indophenol solution. The ascorbic acid was estimated as

$$\text{Ascorbic acid (mg/100 g)} = C \times V \times \text{DF}/W \times 100$$

where C: mg ascorbic acid/mL dye; V: volume of aliquot used for titration; DF: dilution factor; W: weight of sample.

3.10. Microbial Shelf-Life

After baking and cooling, muffins were packed in metallic polyethylene zip lock bag (32-micron) and stored at room temperature (25 °C). The muffins were examined for microbial growth at intervals of 3 days. The microbial shelf life was carried out using the plate count method to determine the total plate counts using Nutrient Agar (HiMedia, Mumbai, India) and yeasts and molds using Potato Dextrose Agar (HiMedia). Potato Dextrose Agar plates were incubated at 25 °C for 48 h and Nutrient Agar plates were incubated at 37 °C for 24 h and the results were expressed as colony-forming units per gram (cfu/g).

3.11. Statistical Analysis

The experiment was statistically analyzed using the Box–Behnken Design based on the RSM. Quadratic ANOVA was used to assess each response variable. ANOVA was used to examine the influence of different variables on the outcome. For comparing the means between the control and the optimized muffin, ANOVA using excel at *p*-value 0.05 significance level was used.

4. Conclusions

The use of the Response Surface Methodology allowed us to optimize the Roselle muffin and achieved significant enhancement of its quality parameters: consumer acceptability, nutritional properties, and shelf-life. Using the BBD model of RSM it was possible to predict the quality parameters of a new type of muffin. The outcome of the study showed that adding Roselle calyx extract to muffins increases their nutritional value by obtaining better antioxidants, bioactive components, total phenolic, and vitamin c that may have the potential of having many health benefits. Microbial shelf-life indicated that Roselle muffin can be stored for 6 days at room temperature with no preservatives added.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/foods11243982/s1>, Figure S1: Normal plot of residual and predicted vs. actual; Table S1: Sensory respondents' score of different muffin formulation resulting from RSM.

Author Contributions: N.K.: Conceptualization, Resources, Supervision, Funding acquisition, Writing—Review and Editing; S.M.: Conceptualization, Methodology, Experimentation, Writing—Original Draft. E.S.: Writing—Review and Editing; E.F.: Funding, Writing—Review; A.D.: Supervision, Writing—Review and Editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by RCN, Norway for Re-FOOD project: International Partnership for Research and Education in energy efficient resource utilization in Food value chains between Norway and India, (Project No. 261709) and Ministry of Tribal Affairs under the scheme "National Fellowship for ST students.

Data Availability Statement: The data presented in this study are available in supplementary material.

Acknowledgments: Shalini Rudra for allowing her lab to undertake texture analysis, V.K. Modi for assisting with sensory analysis, and Asha Kumari for handling the spectrophotometer is greatly acknowledged. We also greatly acknowledged all the panelists involved in the sensory evaluation of the samples.

Conflicts of Interest: The authors declare no conflict of interest or any personal circumstances or interest that may be perceived as inappropriately influencing the representation or interpretation of reported research results.

References

1. Thomford, N.E.; Senthebane, D.A.; Rowe, A.; Munro, D.; Seele, P.; Maroyi, A.; Dzobo, K. Natural products for drug discovery in the 21st century: Innovations for novel drug discovery. *Int. J. Mol. Sci.* **2018**, *19*, 1578. [CrossRef]
2. Kroyer, G.T. Red clover extract as antioxidant active and functional food ingredient. *Innov. Food Sci. Emerg. Technol.* **2004**, *5*, 101–105. [CrossRef]
3. Blunt, J.W.; Carroll, A.R.; Copp, B.R.; Davis, R.A.; Keyzers, R.A.; Prinsep, M.R. Marine natural products. *Nat. Prod. Rep.* **2018**, *35*, 8–53. [CrossRef]
4. Havey, A.L. Natural products in drug discovery. *Drug Discov. Today* **2008**, *13*, 894–901. [CrossRef]
5. Gil-Chávez, G.J.; Villa, J.A.; Ayala-Zavala, J.F.; Heredia, J.B.; Sepulveda, D.; Yahia, E.M.; González-Aguilar, G.A. Technologies for extraction and production of bioactive compounds to be used as nutraceuticals and food ingredients: An overview. *Compr. Rev. Food Sci. Food Saf.* **2013**, *12*, 5–23. [CrossRef]
6. Marak, S.; Shumilina, E.; Kaushik, N.; Falch, E.; Dikiy, A. Effect of Different Drying Methods on the Nutritional Value of *Hibiscus sabdariffa* Calyces as Revealed by NMR Metabolomics. *Molecules* **2021**, *26*, 1675. [CrossRef]
7. Ali, B.H.; Wabel, N.A.; Blunden, G. Phytochemical, pharmacological and toxicological aspects of Hibiscus. *Toxicol. Eval. Nat. Prod. Deriv.* **2005**, *19*, 369–375. [CrossRef]
8. González, C.S.; Balderas, F.T.V.; Regules, A.O.; Beltrán, J.A.G. Antioxidant properties and color of *Hibiscus sabdariffa* extracts. *Cien. Inv. Agr.* **2012**, *39*, 79–90. [CrossRef]
9. Prabawati, N.B.; Oktavirina, V.; Palma, M.; Setyaningsih, W. Edible flowers: Antioxidant compounds and their functional properties. *Horticulturae* **2021**, *7*, 66. [CrossRef]
10. Da-Costa-Rocha, I.; Bonnlaender, B.; Sievers, H.; Pischel, I.; Heinrich, M. *Hibiscus sabdariffa* L.—A phytochemical and pharmacological review. *Food Chem.* **2014**, *165*, 424–443. [CrossRef] [PubMed]
11. Martins, N.; Roriz, C.L.; Morales, P.; Barros, L.; Ferreira, I.C. Food colorants: Challenges, opportunities, and current desires of agro-industries to ensure consumer expectations and regulatory practices. *Trends Food Sci. Technol.* **2016**, *52*, 1–15. [CrossRef]
12. Shruti, V.H.; Ramachandra, C.T.; Nidoni, U.; Hiregoudar, S.; Naik, N.; Kurubar, A.R. Roselle (*Hibiscus sabdariffa* L.) as a source of natural colour: A review. *Plant Arch.* **2016**, *16*, 515–522. Available online: <http://plantarchives.org/PDF%20162/515-522.pdf> (accessed on 3 August 2022).
13. Abou-Arab, A.A.; Abu-Salem, F.M.; Abou-Arab, E.A. Physico-chemical properties of natural pigments (anthocyanin) extracted from Roselle calyces (*Hibiscus sabdariffa*). *J. Am. Sci.* **2011**, *7*, 445–456. Available online: <http://www.americanscience.org/> (accessed on 3 August 2022).
14. Sanz, T.; Salvador, A.; Baixauli, R.; Fiszman, S.M. Evaluation of four types of resistant starch in muffins. II. Effects in texture, colour and consumer response. *Eur. Food Res. Technol.* **2009**, *229*, 197–204. [CrossRef]
15. Martínez-Cervera, S.; Sanz, T.; Salvador, A.; Fiszman, S.M. Rheological, textural, and sensorial properties of low-sucrose muffins reformulated with sucralose/polydextrose. *LWT—Food Sci. Technol.* **2012**, *45*, 213–220. [CrossRef]
16. Liu, R.H. Health-promoting components of fruits and vegetables in the diet. *Adv. Nutr.* **2013**, *4*, 384S–392S. [CrossRef]

17. Rothwell, J.A.; Medina-Remón, A.; Pérez-Jiménez, J.; Neveu, V.; Knaze, V.; Slimani, N.; Scalbert, A. Effects of food processing on polyphenol contents: A systematic analysis using Phenol-Explorer data. *Mol. Nutr. Food Res.* **2015**, *59*, 160–170. [CrossRef]
18. Blessington, T.; Nzaramba, M.N.; Scheuring, D.C.; Hale, A.L.; Reddivari, L.; Miller, J.C. Cooking methods and storage treatments of potato: Effects on carotenoids, antioxidant activity, and phenolics. *Am. J. Potato Res.* **2010**, *87*, 479–491. [CrossRef]
19. Xu, X.; Li, W.; Lu, Z.; Beta, T.; Hydamaka, A.W. Phenolic content, composition, antioxidant activity, and their changes during domestic cooking of potatoes. *J. Agric. Food Chem.* **2009**, *57*, 10231–10238. [CrossRef]
20. Doerry, W.T. *Breadmaking Technology*; American Institute of Baking: New York, NY, USA, 1995; pp. 156–159.
21. Gilbert, R.J.; De Louvois, J.; Donovan, T.; Little, C.; Nye, K.; Ribeiro, C.D.; Bolton, F.J. Guidelines for the microbiological quality of some ready-to-eat foods sampled at the point of sale. PHLS Advisory Committee for Food and Dairy Products. *Commun. Dis. Public Health* **2000**, *3*, 163–167. Available online: <https://www.mb-labs.com/wp-content/uploads/2014/08/Micro-Limits-Ready-to-Eat-Foods.pdf> (accessed on 3 August 2022).
22. Carl, A.B. Confectionery products—Cakes and pastries. In *Encyclopedia of Food Microbiology*; Batt, C.A., Ed.; Academic Press: London, UK, 2014; pp. 500–502.
23. Poliński, S.; Topka, P.; Tańska, M.; Kowalska, S.; Czaplicki, S.; Szydłowska-Czerniak, A. Effect of Grinding Process Parameters and Storage Time on Extraction of Antioxidants from Ginger and Nutmeg. *Molecules* **2022**, *27*, 7395. [CrossRef]
24. Demirok, N.T.; Yıkıncı, S. Combined Effect of Ultrasound and Microwave Power in Tangerine Juice Processing: Bioactive Compounds, Amino Acids, Minerals, and Pathogens. *Processes* **2022**, *10*, 2100. [CrossRef]
25. Barba, F.J.; Rajha, H.N.; Debs, E.; Abi-Khattar, A.M.; Khabbaz, S.; Dar, B.N.; Louka, N. Optimization of Polyphenols' Recovery from Purple Corn Cobs Assisted by Infrared Technology and Use of Extracted Anthocyanins as a Natural Colorant in Pickled Turnip. *Molecules* **2022**, *27*, 5222. [CrossRef] [PubMed]
26. Vo, T.P.; Duong, N.H.N.; Phan, T.H.; Mai, T.P.; Nguyen, D.Q. Optimized Cellulase-Hydrolyzed Deoiled Coconut Cake Powder as Wheat Flour Substitute in Cookies. *Foods* **2022**, *11*, 2709. [CrossRef] [PubMed]
27. Turabi, E.; Sumnu, G.; Sahin, S. Optimization of baking of rice cakes in infrared–microwave combination oven by response surface methodology. *Food Bioprocess Technol.* **2008**, *1*, 64–73. [CrossRef]
28. Sanchez, H.D.; Osella, C.A.; De La Torre, M.A. Use of response surface methodology to optimize gluten-free bread fortified with soy flour and dry milk. *Food Sci. Technol. Int.* **2004**, *10*, 5–9. [CrossRef]
29. Kurek, M.A.; Moczowska-Wyrwisz, M.; Wyrwisz, J.; Karp, S. Development of Gluten-Free Muffins with β -Glucan and Pomegranate Powder Using Response Surface Methodology. *Foods* **2021**, *10*, 2551. [CrossRef]
30. Noordin, M.Y.; Venkatesh, V.C.; Sharif, S.; Elting, S.; Abdullah, A. Application of response surface methodology in describing the performance of coated carbide tools when turning AISI 1045 steel. *J. Mater. Process. Technol.* **2004**, *145*, 46–58. [CrossRef]
31. Siti Faridah, M.A.; Hasmadi, M.; Lee, J.A.; Nurul, S.R. Physicochemical and organoleptic evaluation of muffin partially substituted with roselle calyces (*Hibiscus sabdariffa*) powder. *Borneo Sci.* **2014**, *35*, 10–17. Available online: <http://borneoscience.ums.edu.my/wp-content/uploads/2016/03/physicochemical-and-organoleptic-evaluation-of-muffin-partially-substituted-with-roselle-calyces-hibiscus-sabdariffa-l-powder.pdf> (accessed on 4 August 2022).
32. Everitt, M. Consumer-Targeted Sensory Quality. In *Global Issues in Food Science and Technology*; Academic Press: London, UK, 2009; pp. 117–128. [CrossRef]
33. Krinsky, N.I.; Beecher, G.R.; Burk, R.F.; Chan, A.C.; Erdman, J.J.; Jacob, R.A.; Traber, M.G. Dietary reference intakes for vitamin C, vitamin E, selenium, and carotenoids. *Inst. Med.* **2000**, *19*, 95–185. Available online: <https://www.ncbi.nlm.nih.gov/books/NBK225480/> (accessed on 4 August 2022).
34. Rodríguez-García, J.; Sahi, S.S.; Hernando, I. Functionality of lipase and emulsifiers in low-fat cakes with inulin. *LWT—Food Sci. Technol.* **2014**, *58*, 173–182. [CrossRef]
35. Xu, Q.; Li, B.; Wang, D.; Luo, L.; Liu, G.; Zhou, Y. Microencapsulation and stability analysis of blueberry anthocyanins. In Proceedings of the IOP Conference Series: Earth and Environmental Science, Xi'an, China, 15–16 December 2018; IOP Publishing: Bristol, UK, 2019; Volume 252, p. 052133.
36. Rodriguez, M.V.; Medina, L.M.; Jordano, R. Prolongation of shelf life of sponge cakes using modified atmosphere packaging. *Acta Aliment.* **2002**, *31*, 191–196. [CrossRef]
37. Rafeeda. Simple Vanilla Muffins—The Big Sweet Tooth. 2019. Available online: <https://www.thebigsweettooth.com/simple-vanilla-muffins/> (accessed on 17 July 2021).
38. Horwitz, W. *Official Method of Analysis*, 13th ed.; Association of Analytical Chemists: Washington, DC, USA, 1980.
39. Jauharah, M.A.; Rosli, W.W.; Robert, S.D. Physicochemical and sensorial evaluation of biscuit and muffin incorporated with young corn powder. *Sains Malays.* **2014**, *43*, 45–52. Available online: http://journalarticle.ukm.my/6814/1/06_M.Z._Anis_Jauharah.pdf (accessed on 3 October 2022).
40. Abdel-Moemin, A.R. Effect of Roselle calyces extract on the chemical and sensory properties of functional cupcakes. *Food Sci. Hum. Wellness* **2016**, *5*, 230–237. [CrossRef]
41. Shirazi, O.U.; Khattak, M.M.A.K.; Shukri, N.A.M.; Nasyriq, M.N. Determination of total phenolic, flavonoid content and free radical scavenging activities of common herbs and spices. *J. Pharmacogn. Phytochem.* **2014**, *3*, 104–108. Available online: <https://www.phytojournal.com/archives/2014/vol3issue3/PartB/29.1.pdf> (accessed on 20 August 2022).

42. Lee, J.; Durst, R.W.; Wrolstad, R.E.; Collaborators: Eisele, T.; Giusti, M.M.; Hach, J.; Hofsommer, H.; Koswig, S.; Krueger, D.A.; Kupina, S.; et al. Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: Collaborative study. *J. AOAC Int.* **2005**, *88*, 1269–1278. [[CrossRef](#)]
43. Brad-Williams, W.; Cuvelier, M.E.; Berset, C.L.W.T. Use of a free radical method to evaluate antioxidant activity. *LWT—Food Sci. Technol.* **1995**, *28*, 25–30. [[CrossRef](#)]